

Synthesis of trans-mono(silyl)palladium(II) bromide complexes

Article (Published Version)

Ansell, Melvyn B, Kostakis, George E, Navarro, Oscar and Spencer, John (2021) Synthesis of trans-mono(silyl)palladium(II) bromide complexes. *Molecules*, 26 (9). a2460 1-6. ISSN 1420-3049

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/98642/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Communication

Synthesis of *trans*-Mono(silyl)palladium(II) Bromide Complexes

Melvyn B. Ansell ¹, George E. Kostakis ¹ , Oscar Navarro ^{1,2,*} and John Spencer ^{1,*} 

¹ Department of Chemistry, School of Life Sciences, University of Sussex, Brighton BN1 9QJ, UK; mbansell25@gmail.com (M.B.A.); G.Kostakis@sussex.ac.uk (G.E.K.)

² AMPAC Fine Chemicals, Highway 50 and Hazel Avenue, Building 05-019, Rancho Cordova, CA 95741-1718, USA

* Correspondence: oscar.navarro@apfc.com (O.N.); j.spencer@sussex.ac.uk (J.S.); Tel.: +1-916-357-6928 (O.N.); +44-1273-877-374 (J.S.)

Abstract: The stoichiometric reaction of *cis*-[Pd(ITMe)₂(SiR₃)₂], where (SiR₃ = SiMe₃ and SiMe₂Ph and ITMe = 1,3,4,5-tetramethylimidazol-2-ylidene) with allyl bromide affords the corresponding allylsilanes along with complexes of the type *trans*-[Pd(ITMe)₂(SiR₃)(Br)]. The structure of *trans*-[Pd(ITMe)₂(SiMe₂Ph)Br] **2b** has been determined in the solid state and displays a slightly distorted square-planar geometry with the two N-heterocyclic carbene ligands in a *trans*-configuration.

Keywords: palladium; silicon; N-heterocyclic carbene; allylsilane



Citation: Ansell, M.B.; Kostakis, G.E.; Navarro, O.; Spencer, J. Synthesis of *trans*-Mono(silyl)palladium(II) Bromide Complexes. *Molecules* **2021**, *26*, 2460. <https://doi.org/10.3390/molecules26092460>

Academic Editor: Vincent Ritleng

Received: 13 April 2021

Accepted: 21 April 2021

Published: 23 April 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.




Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Mono(silyl)palladium(II) halide species are purported intermediates in a number of catalytic routes towards allylsilanes [1,2]. Palladium pincer chemistry accounts of such complexes are rather numerous, although examples of their isolation in this catalytic cycle are rare [3–7]. *Trans*-[PdCl(SiF₂Ph)(L)₂] (L = PMe₃, PMe₂Ph or PMePh₂) and allyl bromide were shown to react to afford *trans*-[Pd(L)₂(SiF₂Ph)(Br)] and the corresponding allylsilane [1], and [(^tBuPAR₂)Pd(SiMe₃)(I)] (Ar = 3,5-Me₂-4-OMe-C₆H₂) was synthesized from stoichiometric quantities of [(cod)Pd(CH₂SiMe₃)₂] (cod = 1,5-cyclooctadiene), ^tBuPAR₂ and Me₃SiI [2]. Analogues have been used in silyl-Negishi couplings [8]. We wish to report here our preliminary findings on the reaction of (ITMe)₂Pd(silyl)₂ complexes with allyl bromide (ITMe = 1,3,4,5-tetramethylimidazol-2-ylidene) [9–12].

2. Results and Discussion

The bis(silyl)palladium complexes, *cis*-[Pd(ITMe)₂(SiR₃)₂] (**1a**: SiR₃ = SiMe₃ and **1b**: SiMe₂Ph [13,14] were reacted with excess allylbromide at room temperature under an nitrogen atmosphere to yield *trans*-[Pd(ITMe)₂(SiMe₃)(Br)] **2a** and *trans*-[Pd(ITMe)₂(SiMe₂Ph)Br] **2b** in 92 and 93% yield, respectively (Scheme 1). Reaction progress was monitored by ¹H NMR spectroscopy. Characteristic resonances corresponding to silanes **3a** and **3b** were observed (in a 1:1 stoichiometry with **2a**/**2b**, respectively, upon examination of the crude mixtures).



Scheme 1. Stoichiometric synthesis of mono(silyl)palladium bromide complexes.

In order to further characterize the organometallic complexes, single crystals of **2b** suitable for X-ray analysis were grown by slow evaporation of a saturated deuterated benzene solution at room temperature. X-ray analysis revealed that **2b** displays a marginally distorted square-planar geometry with the two NHCs in a *trans*-configuration and orthogonal to the Br-Pd-Si plane (Figure 1, Table 1).





Figure 1. Molecular structure of **2b**. Hydrogen atoms are omitted for clarity. Selected bond lengths [Å] and angles [°]: Pd1–Br1 2.6333(7), Pd–Si1 2.2948(18), Pd1–C1 2.028(5), Pd1–C8 2.025(5); C1–Pd1–Br1 94.97(16), C1–Pd1–Si1 89.15(17), C8–Pd1–Br1 87.62(16), C8–Pd1–Si1 88.60(17), C1–Pd1–C8 177.2(2).

Table 1. Crystal data and structure refinement for **2b**.

Empirical formula	C ₂₂ H ₃₅ BrN ₄ PdSi
Formula weight	569.94
Temperature/K	173
Crystal system	orthorhombic
Space group	P2 ₁ 2 ₁ 2 ₁
a/Å	10.5467(4)
b/Å	14.3455(3)
c/Å	16.7301(4)
α/°	90
β/°	90
γ/°	90
Volume/Å ³	2531.23(13)
Z	4
ρ _{calc} /cm ³	1.496
μ/mm ^{−1}	2.374
F(000)	1160.0
Crystal size/mm ³	0.22 × 0.2 × 0.15
Radiation	MoKα (λ = 0.71073)
2θ range for data collection/°	6.836 to 52.744
Index ranges	−13 ≤ h ≤ 8, −17 ≤ k ≤ 11, −14 ≤ l ≤ 20
Reflections collected	7612
Independent reflections	4799 [R _{int} = 0.0320, R _{sigma} = 0.0568]
Data/restraints/parameters	4799/0/272
Goodness-of-fit on F ²	1.018
Final R indexes [I ≥ 2σ (I)]	R ₁ = 0.0322, wR ₂ = 0.0614
Final R indexes [all data]	R ₁ = 0.0376, wR ₂ = 0.0640
Largest diff. peak/hole/e Å ^{−3}	0.51/−0.34
Flack parameter	0.004(8)
CCDC deposition number	2076437

The carbenic carbon-Pd bond lengths in **2b** [2.028(5) and 2.025(5) Å] are significantly shorter than in *cis*-[Pd(ITMe)₂(SiMe₂Ph)₂] [2.105(3) and 2.123(3) Å], suggesting SiMe₂Ph exhibits a stronger *trans*-influence than ITMe [15]. The decreased length of the Pd-Si bond in **2b** [2.2948(18) Å] versus *cis*-[Pd(ITMe)₂(SiMe₂Ph)₂] [2.3445(8) and 2.3346(8) Å] infers a stronger Pd-Si bond in **2b** and demonstrates the weak *trans*-influence of Br. Based on these data, the intensity of the *trans*-influence in these two structures follows the sequence: Br < ITMe < SiMe₂Ph. Thus, the preference for the *trans*-configuration observed in **2b** may be attributed to the high *trans*-influence of SiMe₂Ph and the large steric size of Br.

A possible mechanism for the formation of **2** includes either a σ -bond metathesis between a Pd-Si, in *cis*-[Pd(ITMe)₂(SiR₃)₂], and Br-C bond, in allylbromide, or an S_N2/S_N2' by the nucleophilic Pd-Si bond at the electrophilic sites in the allyl halide, leading to a *trans* complex. As we have previously suggested using computational studies on related bis-ITMe complexes, an NHC would then dissociate from the palladium center followed by a *cis* to *trans* isomerization of the Br and Si moieties (Scheme 2) [11]. Finally, the dissociated NHC would re-coordinate, constrained by the bulk of the other ligands, in a *cis*-configuration [16,17].



Scheme 2. Possible mechanistic routes for the formation of **2**.

3. Experimental

The handling of air-sensitive compounds and their spectroscopic measurements were undertaken using standard Schlenk line techniques using pre-dried Ar (using a BASF R3-11(G) catalyst and 4 Å molecular sieves), or in a MBraun glovebox under N₂ (O₂ < 10.0 ppm). All glassware was dried in a 160 °C oven prior to use. Celite was predried in a 200 °C oven and then dried with a heat gun under a dynamic vacuum prior to use. Filter cannulae equipped with microfiber filters were dried in an oven at 160 °C prior to use. Solvents employed in air-sensitive reactions were dried using vacuum distillation, followed by distillation over potassium or stored over activated 4 Å molecular sieves under an Ar atmosphere. NMR spectra were recorded on a Varian VNMRS 400 (Palo Alto, CA, USA) (¹H 399.5 MHz; ¹³C{¹H} 100.5 MHz; ¹¹B{¹H} 128.2 MHz; ¹⁹F 375.9 MHz; ²⁹Si{¹H}

79.4 MHz), or 500 (^1H 499.9 MHz; $^{13}\text{C}\{^1\text{H}\}$ 125.7 MHz). Chemical shifts are reported in ppm. All other experimental details are outlined elsewhere [10].

Synthesis of *trans*-[Pd(ITMe)₂(SiMe₃)(Br)] (**2a**) and Allyltrimethylsilane (**3a**)

Allylbromide (0.032 g, 0.26 mmol) was added to a solution of *cis*-[Pd(ITMe)₂(SiMe₃)₂] (0.043 g, 0.09 mmol) in C₆D₆ or toluene (3.0 mL) and the resulting reaction mixture was stirred at room temperature for 1.5 h. At this stage, the volatiles were removed in vacuo and the off-white powder was washed with hexane (3 × 4.0 mL).

2a, Yield: 0.040 g, 92%. ^1H NMR (399.5 MHz, C₆D₆): δ_{H} = 3.68 [s, 12H, N(1,3)-CH₃], 1.42 [s, 12H, C(4,5)-CH₃], 0.12 [s, 9H, SiMe₃]. $^{13}\text{C}\{^1\text{H}\}$ NMR (100.5 MHz, C₆D₆): δ_{C} = 184.9 [NCN], 124.0 [C(4,5)-CH₃], 35.1 [N(1,3)-CH₃], 8.5 [C(4,5)-CH₃], 6.9 [SiMe₃]. $^{29}\text{Si}\{^1\text{H}\}$ NMR (79.4 MHz, C₆D₆): δ_{Si} = 7.68. Elem. Anal. Calcd. for C₁₇H₃₃N₄SiBrPd: C, 40.20%; H, 6.55%; N, 11.03%. Found: C, 40.15%; H, 6.54%; N, 10.95%. **3a** (from crude reaction solution), ^1H NMR (399.5 MHz, C₆D₆): δ_{H} = 5.77 [m, 1H, CH=], 4.92 [m, 1H, CH=], 4.89 [m, 1H, CH=], 1.44 [m, 2H, CH₂], −0.03 [s, 9H, SiMe₃]. [Agrees with an independently taken ^1H NMR sample of commercially available allyltrimethylsilane].

Synthesis of *trans*-[Pd(ITMe)₂(SiMe₂Ph)(Br)] (**2b**)

Allylbromide (6.0 μL , 0.07 mmol) and *cis*-[Pd(ITMe)₂(SiMe₂Ph)] (0.021 g, 0.03 mmol) were dissolved in C₆D₆ or toluene (1.0 mL). The resulting reaction mixture was stirred at room temperature for 2 h under an N₂ atmosphere. At this stage, all volatiles were removed in vacuo and the resulting white solid was washed with hexane (3 × 2.0 mL). Yield: 0.018 g, 93%. ^1H NMR (399.5 MHz, C₆D₆): δ_{H} = 7.20 [m, 2H, SiMe₂Ph], 7.07 [m, 3H, SiMe₂Ph], 3.51 [s, 12H, N(1,3)-CH₃], 1.42 [s, 12H, C(4,5)-CH₃], 0.31 [s, 6H, SiMe₂Ph]. $^{13}\text{C}\{^1\text{H}\}$ NMR (100.5 MHz, C₆D₆): δ_{C} = 183.4 [NCN], 149.6 [SiMe₂*i*-Ph], 133.1 [SiMe₂Ph], 127.0 [SiMe₂Ph], 126.5 [SiMe₂*p*-Ph], 124.2 [C(4,5)-CH₃], 34.9 [N(1,3)-CH₃], 8.5 [C(4,5)-CH₃], 4.2 [SiMe₂Ph]. $^{29}\text{Si}\{^1\text{H}\}$ NMR (79.4 MHz, C₆D₆): δ_{Si} = 2.44. (It was not possible to obtain elemental analysis for **2b**—every attempt resulted in numbers that were inconsistent with calculated values. A possible reason for this is decomposition of **2b** by exposure to air or moisture on transit to data collection).

Crystal data for **2b**: C₂₂H₃₅N₄SiBrPd, M_r = 569.94 g mol^{−1}, orthorhombic, space group P2 = 2₁, a = 10.5467(4) Å, b = 14.3455(3) Å, c = 16.7301(4) Å, α = 90°, β = 90°, γ = 90°, V = 2531.23(13) Å³, Z = 4, T = 173 K, $\lambda\text{Mo(K}\alpha)$ = 0.71073, R_1 [$I > 2\sigma(I)$] = 0.0345, wR_2 (all data) = 0.0677, GooF = 1.011.

Crude ^1H NMR data are consistent with the formation of allyldimethylphenylsilane (**3b**) as a product of this reaction. However, this was not isolated in this instance [18].

4. Conclusions

Under mild conditions, non-pincer bis(NHC)(silyl)palladium halide complexes of the type *trans*-[Pd(ITMe)₂(SiR₃)(Br)] (SiR₃ = SiMe₂Ph (**2a**), and SiMe₃ (**2b**)) were synthesized, by the reaction of allylbromide with the corresponding complexes *cis*-[Pd(ITMe)₂(SiR₃)₂], **1a** or **1b**, respectively. A possible mechanistic route for the formation of **2** involves either a σ -bond metathesis or an S_N2/S_N2' reaction between allylbromide and **1**. This would necessitate a *cis-trans* isomerization via dissociation of an NHC ligand-[19]. The reactivity of *trans*-[Pd(ITMe)₂(SiR₃)(Br)] is unexplored but will soon be carried out. The facile formation and apparent stability of *trans*-**2** may indeed hinder the catalytic silylation of allyl halides mediated by ITMe₂Pd-based complexes since the adoption of a *cis*-configuration is a prerequisite for reductive elimination and involvement in a catalytic cycle. Solutions to these unexplored questions are currently being sought, e.g., the potential for halide abstraction, and will be reported in due course.

Author Contributions: Conceptualization, M.B.A., O.N., J.S. Writing—original draft preparation, all authors; writing—review and editing, all authors. X-ray data acquisition and refinement: M.B.A. and G.E.K. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by an EPSRC Standard Research Studentship (DTG, Grant # EP/L505109/1), to M.B.A.

Institutional Review Board Statement: Not applicable.

Acknowledgments: Dedicated to Michel Pfeffer for his contributions to C-H activation, palladium, and ruthenium chemistry. A gentleman and a scholar.

Conflicts of Interest: The authors declare no conflict of interest.

Sample Availability: No samples of the compounds are available from the authors.

Disclosure: This work has been previously presented as part of a thesis available at: Ansell, Melvyn B (2017) *Novel (N-heterocyclic carbene)-palladium(0) complexes as catalysts in element-element bond additions to unsaturated moieties*. Doctoral thesis (PhD), University of Sussex, Brighton BN1 9QJ, UK; <http://sro.sussex.ac.uk/id/eprint/68072/>, accessed on 12 May 2017).

References

- Ozawa, F.; Sugawara, M.; Hasebe, K.; Hayashi, T. Reactions of bis(silyl)palladium(II) complexes with allyl halides. Synthesis of mono(silyl)palladium(II) halides and X-ray structure of trans-PdCl(SiF₂Ph)(PMe₂Ph)₂. *Inorg. Chim. Acta* **1999**, *296*, 19–25. [CrossRef]
- McAtee, J.R.; Yap, G.P.A.; Watson, D.A. Rational design of a second generation catalyst for preparation of allylsilanes using the silyl-Heck reaction. *J. Am. Chem. Soc.* **2014**, *136*, 10166–10172. [CrossRef] [PubMed]
- MacInnis, M.C.; MacLean, D.F.; Lundgren, R.J.; McDonald, R.; Turculet, L. Synthesis and reactivity of platinum group metal complexes featuring the new pincer-like bis (phosphino) silyl ligand [K₃-(2-Ph 2PC₆H₄)₂SiMe]₃-([PSiP]): Application in the ruthenium-mediated transfer hydrogenation of ketones. *Organometallics* **2007**, *26*, 6522–6525. [CrossRef]
- Takaya, J.; Nakamura, S.; Iwasawa, N. Synthesis, structure, and catalytic activity of palladium complexes bearing a tridentate pcp-pincer ligand of heavier group 14 element (X = Ge, Sn). *Chem. Lett.* **2012**, *41*, 967–969. [CrossRef]
- Ruddy, A.J.; Mitton, S.J.; McDonald, R.; Turculet, L. Hemilabile silyl pincer ligation: Platinum group PSiN complexes and triple C–H activation to form a (PSiC)Ru carbene complex. *Chem. Commun.* **2012**, *48*, 1159–1161. [CrossRef]
- Wahlisch, S.; Brendler, E.; Heine, T.; Zhechkov, L.; Wagler, J. 7-Azaindol-1-yl(organo)silanes and Their PdCl₂ Complexes: Pd-Capped Tetrahedral Silicon Coordination Spheres and Paddlewheels with a Pd–Si Axis. *Organometallics* **2014**, *33*, 2479–2488. [CrossRef]
- Suh, H.W.; Balcells, D.; Edwards, A.J.; Guard, L.M.; Hazari, N.; Mader, E.A.; Mercado, B.Q.; Repisky, M. Understanding the Solution and Solid-State Structures of Pd and Pt PSiP Pincer-Supported Hydrides. *Inorg. Chem.* **2015**, *54*, 11411–11422. [CrossRef] [PubMed]
- Cinderella, A.P.; Vulovic, B.; Watson, D.A. Palladium-Catalyzed Cross-Coupling of Silyl Electrophiles with Alkylzinc Halides: A Silyl-Negishi Reaction. *J. Am. Chem. Soc.* **2017**, *139*, 7741–7744. [CrossRef] [PubMed]
- Ansell, M.B.; Furfari, S.K.; Cloke, F.G.N.; Roe, S.M.; Spencer, J.; Navarro, O. Comparison of the Reactivity of the Low Buried-Volume Carbene Complexes (ITMe)₂Pd(PhC≡CPh) and (ITMe)₂Pd(PhN=NPh). *Organometallics* **2018**, *37*, 1214–1218. [CrossRef]
- Ansell, M.B.; Menezes Da Silva, V.H.; Heerdt, G.; Braga, A.A.C.; Spencer, J.; Navarro, O. An experimental and theoretical study into the facile, homogenous (N-heterocyclic carbene)₂-Pd(0) catalyzed diboration of internal and terminal alkynes. *Catal. Sci. Technol.* **2016**, *6*, 7461–7467. [CrossRef]
- Ansell, M.B.; Kostakis, G.E.; Braunschweig, H.; Navarro, O.; Spencer, J. Synthesis of Functionalized Hydrazines: Facile Homogeneous (N-Heterocyclic Carbene)-Palladium(0)-Catalyzed Diboration and Silaboration of Azobenzenes. *Adv. Synth. Catal.* **2016**, *358*. [CrossRef] [PubMed]
- Ansell, M.B.; Navarro, O.; Spencer, J. Transition metal catalyzed element–element additions to alkynes. *Coord. Chem. Rev.* **2017**, *336*. [CrossRef]
- Ansell, M.B.; Roberts, D.E.; Cloke, F.G.N.; Navarro, O.; Spencer, J. Synthesis of an [(NHC)₂Pd(SiMe₃)₂] complex and catalytic cis-bis(silyl)ations of alkynes with unactivated disilanes. *Angew. Chem. Int. Ed.* **2015**, *54*, 5579–5582. [CrossRef] [PubMed]
- Ansell, M.B.; Spencer, J.; Navarro, O. (N-Heterocyclic Carbene)₂-Pd(0)-Catalyzed Silaboration of Internal and Terminal Alkynes: Scope and Mechanistic Studies. *ACS Catal.* **2016**, *6*, 2192–2196. [CrossRef]
- Mo, Z.; Liu, Y.; Deng, L. Anchoring of silyl donors on a N-heterocyclic carbene through the cobalt-mediated silylation of benzylic C–H bonds. *Angew. Chem. Int. Ed.* **2013**, *52*, 10845–10849. [CrossRef] [PubMed]
- De, K.; Lewis, A.K.; Caddick, S.; Cloke, F.G.N.; Billingham, N.C.; Hitchcock, P.B.; Leonard, J. Synthetic, structural, and mechanistic studies on the oxidative addition of aromatic chlorides to a palladium (N-heterocyclic carbene) complex: Relevance to catalytic amination. *J. Am. Chem. Soc.* **2003**, *125*, 10066–10073.
- De, K.; Lewis, A.K.; Caddick, S.; Esposito, O.; Cloke, F.G.N.; Hitchcock, P.B. Synthetic and structural studies on amine coordination to Pd-N-heterocyclic carbene complexes. *Dalt. Trans.* **2009**, *35*, 7094–7098.

-
18. Godineau, E.; Schenk, K.; Landais, Y. Synthesis of fused piperidinones through free-radical-ionic cascade. *J. Org. Chem.* **2008**, *73*, 6983–6993. [[CrossRef](#)] [[PubMed](#)]
 19. Suarez, A.G.; Nelson, D.J.; Nolan, S.P. Quantifying and understanding the steric properties of N-heterocyclic carbens. *Chem. Commun.* **2017**, *53*, 2650–2660. [[CrossRef](#)] [[PubMed](#)]